

# Gene Expression Variation Resolves Species and Individual Strains among Coral-Associated Dinoflagellates within the Genus *Symbiodinium*

John E. Parkinson<sup>1,\*</sup>, Sebastian Baumgarten<sup>2</sup>, Craig T. Michell<sup>2</sup>, Iliana B. Baums<sup>1</sup>, Todd C. LaJeunesse<sup>1</sup>, and Christian R. Voelstra<sup>2,\*</sup>

<sup>1</sup>Department of Biology, Pennsylvania State University

<sup>2</sup>Red Sea Research Center, Division of Biological and Environmental Science and Engineering, King Abdullah University of Science and Technology (KAUST), Thuwal, Saudi Arabia

\*Corresponding author: E-mail: jparkinson@psu.edu; christian.voelstra@kaust.edu.sa.

Accepted: February 1, 2016

**Data deposition:** This project has been deposited at the NCBI Sequence Read Archive database under accession numbers PRJNA274852, PRJNA274854, PRJNA274855, and PRJNA274856, and the RNAseq transcriptomes at reefgenomics.org.

## Abstract

Reef-building corals depend on symbiotic mutualisms with photosynthetic dinoflagellates in the genus *Symbiodinium*. This large microalgal group comprises many highly divergent lineages (“Clades A–I”) and hundreds of undescribed species. Given their ecological importance, efforts have turned to genomic approaches to characterize the functional ecology of *Symbiodinium*. To date, investigators have only compared gene expression between representatives from separate clades—the equivalent of contrasting genera or families in other dinoflagellate groups—making it impossible to distinguish between clade-level and species-level functional differences. Here, we examined the transcriptomes of four species within one *Symbiodinium* clade (Clade B) at ~20,000 orthologous genes, as well as multiple isoclonal cell lines within species (i.e., cultured strains). These species span two major adaptive radiations within Clade B, each encompassing both host-specialized and ecologically cryptic taxa. Species-specific expression differences were consistently enriched for photosynthesis-related genes, likely reflecting selection pressures driving niche diversification. Transcriptional variation among strains involved fatty acid metabolism and biosynthesis pathways. Such differences among individuals are potentially a major source of physiological variation, contributing to the functional diversity of coral holobionts composed of unique host–symbiont genotype pairings. Our findings expand the genomic resources available for this important symbiont group and emphasize the power of comparative transcriptomics as a method for studying speciation processes and interindividual variation in nonmodel organisms.

**Key words:** dinoflagellates, phylogenetics, RNAseq, symbiosis, transcriptome, zooxanthellae.

## Introduction

The concept that adaptation and speciation are driven largely by natural selection on variant individuals of a population is central to evolutionary biology. Much like other types of genetic diversity, gene expression variation is extensive, highly heritable, and subject to selection (Ranz and Machado 2006; Voelstra et al. 2007; Wittkopp et al. 2008). The role of differential gene expression in ecological speciation has received renewed interest in the genomics era because the molecular biology of nonmodel organisms with unique evolutionary histories can now be studied in great detail at relatively low cost (Wolf et al. 2010). For example, among two recently diverged

species of cordgrass, only one is successful at invading environments perturbed by climate change, and it exhibits unique expression patterns at growth- and stress-related genes (Chelaifa et al. 2010). A similar study in daisies illustrated that a comparative transcriptomic framework can be used to identify selective processes affecting ecological speciation (Chapman et al. 2013). Additionally, transcription-based assays of microbial metagenomes have revealed unique niche diversification (e.g., specialization for certain substrates, metabolic pathways, or environments) that is otherwise hidden due to functional redundancy in the genomes of

© The Author 2016. Published by Oxford University Press on behalf of the Society for Molecular Biology and Evolution.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com

many bacteria (Gifford et al. 2013). Thus, comparative genomics can also provide a means to recognize important functional variation in organisms that are difficult to probe phenotypically, such as corals and their symbionts.

Coral reef ecosystems support tremendous marine biodiversity and ecological goods and services (Moberg and Folke 1999). Coral productivity and growth depend on a mutualism with endosymbiotic dinoflagellates known as *Symbiodinium* (Muscatine and Porter 1977; Muscatine 1990; Yellowlees et al. 2008). This microalgal “genus” is incredibly diverse, encompassing at least nine major lineages that show ribosomal divergence equivalent to that found among different genera, families, or even orders of other dinoflagellates (Rowan and Powers 1992). Likewise, *Symbiodinium* exhibit many unique ecologies, ranging from “host-specialized” taxa commonly found as symbiotic partners of corals (Parkinson, Coffroth, et al. 2015), to “ecologically cryptic” taxa with alternate non-symbiotic lifestyles (LaJeunesse et al. 2015), to completely “free-living” taxa that thrive independently in the water column (Jeong et al. 2014). Unlike their mostly obligate coral hosts, many *Symbiodinium* can survive ex hospite and are maintained in culture. In the natural environment, stressful conditions can cause the association between corals and host-specialized symbionts to break down in a process called coral bleaching, which can lead to colony mortality (Fitt et al. 2001). Climate change is predicted to drive more frequent and intense bleaching events (Hoegh-Guldberg 1999), prompting a major research focus on how climate-related stressors might affect coral–dinoflagellate symbioses in the future. Accordingly, the last decade has generated many studies describing coral host transcription in various contexts (Meyer and Weis 2012), but comparable studies in *Symbiodinium* are still in their early stages (Leggat et al. 2007; Leggat, Yellowlees, et al. 2011; Lin 2011).

With the incorporation of next-generation sequencing technology, genomic resources for *Symbiodinium* have expanded greatly despite their status as a nonmodel organism. The first draft genome was released in 2013 (Shoguchi et al. 2013), with the complete chloroplast genome following shortly thereafter (Barbrook et al. 2014). Multiple mRNA transcriptomes are available (Bayer et al. 2012; Ladner et al. 2012; Baumgarten et al. 2013; Rosic et al. 2014; Xiang et al. 2015), representing the four major clades known to associate with scleractinian corals (Clades A, B, C, and D). Recent efforts have expanded in important new directions, such as the description of *Symbiodinium* microRNAs (Baumgarten et al. 2013), the comparison of orthologous genes among clades (Voolstra et al. 2009; Ladner et al. 2012; Barshis et al. 2014; Rosic et al. 2014), the completion of another draft genome (Lin et al. 2015), and the development of the *Aiptasia*–*Symbiodinium* system for in-depth cellular and physiological research (Weis et al. 2008; Sunagawa et al. 2009; Lehnert et al. 2012, 2014; Xiang et al. 2013; Baumgarten et al. 2015).

Dinoflagellate genomes are unique among eukaryotes for multiple reasons (Leggat, Yellowlees, et al. 2011). Of particular

note, dinoflagellates including *Symbiodinium* modulate nuclear-encoded protein levels predominantly by posttranscriptional processes (Morse et al. 1989; Leggat, Seneca, et al. 2011). It is now understood that dinoflagellates also exhibit some measure of transcriptional regulation, albeit changes in expression profiles are minimal when exposed to different environmental conditions (Erdner and Anderson 2006; Moustafa et al. 2010). For example, the number and magnitude of expression changes among *Symbiodinium* exposed to thermal stress are relatively small compared with their animal hosts (Leggat, Seneca, et al. 2011). Barshis et al. (2014) found that two *Symbiodinium* spp. in Clades C and D did not alter gene expression when exposed to temperature stress in hospite, even though the host response involved the modulation of hundreds of genes (Barshis et al. 2013, 2014). Interestingly, a large number of transcriptional differences were maintained, or fixed, for the two species from different clades regardless of temperature treatment (Barshis et al. 2014). This suggests that fixed expression differences are likely to be evident in strains cultured ex hospite under identical controlled environmental conditions. Differences in these “stable-state” expression profiles among lineages may strongly reflect evolutionary divergence, some of which may be adaptive. These expression patterns may also correspond to functional differences among distantly related species. If lineage-specific expression extends to the subcladal level—that is, between species within clades or among individual strains within species—it will be critical to recognize this source of variation when interpreting *Symbiodinium* genomic data and account for it in future experimental designs.

By comparing different isoclonal cell lines (strains), it is possible to reveal intraspecific variation in genomic features that underlie ecological and physiological phenotypes. For example, unique genes distinguish strains of nitrogen-fixing rhizobial bacteria with different symbiotic efficiencies and host specificities (Galardini et al. 2011; Österman et al. 2015). At the level of transcription, toxic and nontoxic strains of the dinoflagellate *Alexandrium minutum* maintain fixed expression differences at shared genes (Yang et al. 2010). We may expect similar patterns among *Symbiodinium* strains, but this idea has never been tested. *Symbiodinium* belonging to Clade B are ideal candidates for further genomic characterization because several ecologically distinct species within this group were recently described (LaJeunesse et al. 2012; Parkinson, Coffroth, et al. 2015), a draft genome exists for the member species *Symbiodinium minutum* (Shoguchi et al. 2013), and multiple genetically distinct cultures are available for several species.

Currently, the extent of variation among species within a single *Symbiodinium* clade and among individual strains within a single species is mostly unknown (Parkinson and Baums 2014). To address this knowledge gap, we analyzed stable-state gene expression among four species representing the two major evolutionary radiations within Clade B: The

Pleistocene (B1) radiation and the Pliocene (B19) radiation (sensu LaJeunesse 2005). For each radiation, two species with different ecologies were studied: Either host-specialized taxa or ecologically cryptic taxa. Although these latter species were originally cultured from coral tissues, they have never been detected as the numerically dominant symbionts in cnidarian mutualisms, and therefore were probably commensals or free-living contaminants isolated from the mucus or gastrovascular cavity (Parkinson, Coffroth, et al. 2015). Where available, we incorporated biological replication in the form of distinct isoclonal cell cultures. The genomic resources developed here should assist in the design and interpretation of future comparative transcriptional analyses among *Symbiodinium* strains, species, and clades, as well as broaden our understanding of speciation among microeukaryotes.

## Materials and Methods

### Culturing

Isoclonal cultures (strains) of Clade B *Symbiodinium* were maintained at the Pennsylvania State University. They were originally acquired from the Robert K. Trench and Buffalo Undersea Reef Research collections. This study included one strain of *Symbiodinium aenigmaticum* (mac04-487), four strains of *S. minutum* (mac703, Mf1.05b, rt002, and rt351), one strain of *Symbiodinium pseudominutum* (rt146), and four strains of *Symbiodinium psymgophilum* (HIAp, Mf10.14b.02, PurPFlex, and rt141), for the analysis of ten individual transcriptomes. Most strains are available from the Provasoli-Guillard National Center for Marine Algae and Microbiota at Bigelow Laboratory for Ocean Sciences, East Boothbay, Maine (LaJeunesse et al. 2012; Parkinson, Coffroth, et al. 2015) or by request. Within *S. minutum* and *S. psymgophilum*, strains were confirmed to represent distinct genotypes based on repeat length variation at the microsatellite locus Sym15 (Pettay and LaJeunesse 2007) and haplotype differences in the *psbA* noncoding region (LaJeunesse and Thornhill 2011).

Single cells were originally isolated from host tissues by Schoenberg and Trench (1980) using modified methods of McLaughlin and Zahl (1959) or by Mary Alice Coffroth following the methods of Santos et al. (2001). To establish initial crude cultures, several drops of a heavy suspension of symbiont cells were transferred into nutrient-enriched filtered seawater (Provasoli 1968) and then spread onto semisolid agar (0.8%) containing the same seawater. Vegetative cells from viable colonies on agar were then transferred to liquid medium ASP-8A (Ahles 1967). To generate isoclonal lines, only individual motile cells were transferred to fresh media. For this experiment, an additional transfer to new media was made to synchronize all cultures. Final cultures were grown in 50 ml volumes in Erlenmeyer flasks for 2 weeks up to concentrations  $\sim 1 \times 10^6$  cells  $\cdot$  ml $^{-1}$ . Cultures were maintained in incubators at 26 °C under Philips fluorescent tubes

(Koninklijke Philips Electronics, Amsterdam, The Netherlands) delivering 80–120  $\mu$ mol  $\cdot$  m $^{-2}$   $\cdot$  s $^{-1}$  photosynthetically active radiation on a 12:12 (light:dark) photoperiod. All cultures grew together under identical conditions until processed simultaneously for RNA extraction.

### RNA Isolation and Sequencing

At the sixth hour of the light photoperiod on the last day of the second week of growth postsynchronization, all target cultures were transferred to 50 ml tubes and centrifuged at 3000 RCF (relative centrifugal force). The media was decanted and the pellets were flash frozen in liquid nitrogen. Pellets were ground with a prechilled mortar and pestle and transferred into 1.5 ml tubes. Nucleic acids were extracted with TriReagent (Thermo Fisher Scientific, Waltham, MA) and RNA was isolated and cleaned with the RNeasy Kit (Qiagen, Venlo, The Netherlands) according to the manufacturer's protocols.

Total RNA isolations were shipped on dry ice to the KAUST Red Sea Research Center, where they were quality checked using a Bioanalyzer 2000 (Agilent, Santa Clara, CA) prior to library preparation. For Illumina 2  $\times$  100 bp paired-end sequencing, 180 bp libraries were generated from oligo(dT)-enriched mRNA using the NEBNext Ultra Directional RNA Library Prep Kit for Illumina (New England Biolabs, Ipswich, MA) according to the manufacturer's protocols. Each read pair ideally yielded a partially overlapping 180 bp contiguous sequence, allowing for additional quality control. mRNA sequencing libraries for each of the ten samples were multiplexed in equimolar concentrations and run on one lane on the Illumina HiSeq 2000 platform, producing a total of 142 million paired-end reads. All raw RNAseq data are available in the NCBI Sequence Read Archive database under accession numbers <http://www.ncbi.nlm.nih.gov/bioproject/PRJNA274856/> (*S. aenigmaticum*), <http://www.ncbi.nlm.nih.gov/bioproject/?term=PRJNA274852> (*S. minutum*), <http://www.ncbi.nlm.nih.gov/bioproject/?term=PRJNA274855> (*S. pseudominutum*), and <http://www.ncbi.nlm.nih.gov/bioproject/?term=PRJNA274854> (*S. psymgophilum*).

### Transcriptome Assemblies and Annotation

Adapters and low quality nucleotides (<20 Phred score in ASCII 33 format) were removed from raw reads with Trimmomatic (Bolger et al. 2014). Reads were error corrected with the error correction module of the AIPATHS-LG pipeline (Gnerre et al. 2011; Ribeiro et al. 2012). Quality-controlled reads combined from all samples on a per-species basis were assembled using the Trinity package (Grabherr et al. 2011) with minimum *k*-mer coverage of 2 and minimum contig length of 250 bp to generate one reference assembly per species (four total). For mapping purposes, we further reduced each reference assembly to include only the longest transcript for a related set of splice variants of a gene. For each sample, reads were mapped back to the reduced assembly for the appropriate species with Bowtie2 (Langmead and

Salzberg 2012), and quantified by summing counts of all transcripts per gene with the program eXpress (Roberts and Pachter 2013), producing effective read counts and FPKM values (Fragments Per Kilobase of transcript per Million mapped reads).

Each reference assembly was annotated by iterative searches of the longest transcript per gene against SwissProt, TrEMBL, and NCBI nr sequence databases (UniProt Consortium 2013; Pruitt et al. 2014) using BLASTX (Altschul et al. 1990) and the October 2013 releases. Only hits with  $e$ -values  $< 1 \times 10^{-5}$  were retained. All genes remaining unannotated after BLASTX against the first database were passed onto the next sequentially. Gene Ontology (GO) categories were assigned through the BLASTX hit to SwissProt or TrEMBL databases, and subsequent mapping to the UniProt-GOA database (Dimmer et al. 2012). The assembled and annotated transcriptome sequences for each species are available at <http://reefgenomics.org>.

The recently published *S. minutum* draft genome (Shoguchi et al. 2013) was derived from strain Mf1.05b, which was included in this study. To compare our sequencing results to this resource, we aligned our Mf1.05b reads to the exome of the draft genome to estimate the proportion of mappable reads using Bowtie 2; 73% of the raw reads aligned.

To assess how comparable our Clade B species assemblies were in terms of gene content independent of expression, complete assemblies were uploaded into Ingenuity Pathway Analysis (IPA) software (Ingenuity Systems, [www.ingenuity.com](http://www.ingenuity.com), last accessed January 2014). IPA compares user-provided gene lists with reference canonical pathways in the manually curated Ingenuity Knowledge Base. It generates a ratio of genes present versus total genes belonging to a pathway and tests for the probability of significant enrichment for that pathway in the *Symbiodinium* transcriptome with Fisher's exact tests. The Ingenuity Knowledge Base is designed mainly for model organism data, so results should be interpreted in the context of pathways that are well annotated and highly conserved across eukaryotes. We were less concerned about pathway identity and more interested in whether representation within a pathway was similar across Clade B species. Thus, we compared ratio values for all transcriptomes at all pathways determined to be significantly enriched in the dataset at  $P_{adj} < 0.05$ . As an additional metric of comparison across Clade B species, we mined all transcriptomes for repeats and flanking regions using the software SciRoKo (Kofler et al. 2007) and considered repeat motifs between 2 and 6 bp.

#### Ortholog Identification and Differential Expression between Species, Lineages, and Ecologies

To test for differential expression between the four species, it was necessary to identify a set of comparable orthologous genes. Open reading frames (ORFs) were predicted for each transcriptome with TransDecoder (Haas et al. 2013).

Orthologous genes were identified via reciprocal BLASTP of ORFs pairwise for each species within the program InParanoid (Remm et al. 2001), retaining hits with bitscores  $> 300$  (supplementary table S1, Supplementary Material online). Multiparanoind (Alexeyenko et al. 2006) was then used to identify orthologs that occurred exactly four times (once in each species). Additional domain-based functional annotations were assigned using the Pfam database v27 (Finn et al. 2014) and are provided in supplementary table S2, Supplementary Material online.

Most current software designed to analyze differential expression for RNAseq data assumes raw read counts among samples mapped to one common reference transcriptome, and therefore only accepts integer values as input. To compare expression at orthologous genes across species, it was necessary to normalize read counts by transcript length using FPKM to account for species-specific sequence length differences. This normalization produced many decimal expression values that were still informative. For this reason, we scaled FPKM by a common factor such that the lowest expressed gene's value equaled 1 and then rounded values to the nearest integer. Thus, a scaled FPKM of 50 means the gene is expressed 50 times higher than the lowest expressed gene retained in the data set. This way, all orthologs could be compared in the scaled FPKM space.

Scaled FPKM data were then used as input for the R package edgeR (Robinson et al. 2010), which accommodates data sets with unequal replication when performing comparisons among treatments (in this case, species). No additional normalization procedures were carried out within the program. Significant differential expression was determined by pairwise comparisons among species with a false discovery rate-adjusted  $P$  value (FDR) of  $< 0.1$ . To additionally test by lineage, all *S. minutum* and *S. pseudominutum* samples were grouped as "Pleistocene" and all *S. aenigmaticum* and *S. psygmophilum* samples were grouped as "Pliocene." To test by ecology, all *S. minutum* and *S. psygmophilum* samples were grouped as host-specialized, and all *S. aenigmaticum* and *S. pseudominutum* samples were grouped as ecologically cryptic.

Multidimensional scaling (MDS) plots were generated in edgeR using the plotMDS function. The distances between pairs of RNA samples correspond to the leading log<sub>2</sub>-fold-changes, which is the average (root-mean-square) of the largest absolute log<sub>2</sub>-fold-changes (Robinson et al. 2010). In all three comparisons of the MDS plots (non-DEG [differentially expressed gene] only, non-DEG + DEG, and DEG only), similar clustering was observed among the four replicates of the two species with replicates. It was therefore reasonable to assume that (hypothetical) replicates of the other two species would show a similar distribution. Because variation between replicates was consistent, we assumed that the distances between any of the species (which contributes to the number of DEGs identified) was not affected by the number of replicates within a species. This was incorporated into the between-species

comparison by calculating the distribution with edgeR once over all samples/replicates before comparing species. One distribution value was used for all subsequent comparisons, including those with species that had only one replicate.

### Differential Expression within Species

For both *S. minutum* and *S. psugmophilum*, isoclinal cultures of four individual strains each were available, providing two opportunities to test for differential expression within a species. Each species was analyzed separately in edgeR (Robinson et al. 2010). Raw read counts were normalized with the geometric mean method.

Unlike hybridization-based techniques, the level of technical variation from sequencing is predictable and can therefore be distinguished from biological variation (Chen et al. 2014). In this case, the technical coefficient of variation describes the measurement error derived from the uncertainty with which the abundance of every gene is estimated from the sequencing platform, which decreases with increasing total counts for each gene in an RNA sample. In contrast, the biological coefficient of variation (BCV) describes the variation of the unknown, true abundance of each gene among replicates of RNA samples that will remain, even if sequencing depth could be increased indefinitely. Thus, the BCV represents the most important and main source of variation in expression studies using a high-throughput deep-sequencing approach (McCarthy et al. 2012). In RNA expression studies, the BCV is usually determined from the biological replicates of RNA samples so the total variation of gene abundances can be calculated by considering the following equation:  $\text{Total CV}^2 = \text{Technical CV}^2 + \text{Biological CV}^2$  (McCarthy et al. 2012).

Due to the lack of replicates for the comparison of the four genotypes of *S. minutum* and *S. psugmophilum*, we set the BCV to a fixed value a priori under the assumption that a majority of genes were not differentially expressed, which we considered appropriate for *Symbiodinium* of the same species under stable-state conditions. Although a value of ~1% is suggested for technical replicates and a value of ~10% is suggested for unique samples from separate but genetically identical model organisms, a value of ~40% is appropriate for independent biological samples (Chen et al. 2014) and was chosen for our expression analysis among the four genotypes of both *Symbiodinium* species.

Significance of DEGs was determined by pairwise comparisons among individuals based on the negative binomial distribution with  $\text{FDR} < 0.1$ . A pairwise Euclidian distance matrix for all strain comparisons within and between *S. minutum* and *S. psugmophilum* was computed based on scaled FPKM values using PRIMER v6 software (Clarke and Gorley 2006).

### Visualization and Functional Analyses

DEGs between and within species were visualized as heatmaps by plotting scaled FPKM expression values with Gene-

E (Gould 2015). Lists were tested for GO term functional enrichment with the R/Bioconductor package topGO (Alexa and Rahnenfuhrer 2010), using the default "weight01" Alexa algorithm with the recommended cutoff of  $P < 0.05$ .

## Results

### Transcriptome Assemblies

We targeted isoclinal strains from four Clade B *Symbiodinium* species: *S. aenigmaticum* ( $n = 1$  strain: mac04-487); *S. minutum* ( $n = 4$  strains: mac703, Mf1.05b, rt002, and rt351); *S. pseudominutum* ( $n = 1$  strain: rt146); and *S. psugmophilum* ( $n = 4$  strains: HIAp, Mf10.14b.02, PurPFlex, and rt141). We reared the ten cultures under identical conditions in one incubator to assess stable-state conditions in the absence of environmental variability. *Symbiodinium minutum* and *S. pseudominutum* belong to the Pleistocene (B1) radiation. The former is a host-specialized mutualist because it commonly associates with the anemone *Aiptasia* sp. The latter is ecologically cryptic—although it has been isolated from the background symbiont population of four cnidarians, it has never been identified as a dominant symbiont. *Symbiodinium psugmophilum* and *S. aenigmaticum* belong to the Pliocene (B19) radiation; the former is host-specialized, the latter is ecologically cryptic. Note the uneven distribution of strains within species. This limitation was based on which cultures were available in the collection and meant that certain contrasts (e.g., 4 *S. minutum* strains vs. 4 *S. psugmophilum* strains) potentially had more power to detect differential expression than others (e.g., 1 *S. aenigmaticum* strain vs. 1 *S. pseudominutum* strain). However, for these data it was unlikely that the number of replicates affected power given that the same variance distribution was used for all comparisons (see Materials and Methods).

From these cultures, we generated ten high-quality short-read RNAseq libraries (table 1). Across species, sequencing statistics were quite similar. Total reads per sample averaged 14.3 million, while on average 88.5% of reads per sample passed quality control. For each of the four species, we generated a single reference assembly from either a combination of all strains of a given species (in the cases of *S. minutum* and *S. psugmophilum*) or from the single representative strain (in the cases of *S. aenigmaticum* and *S. pseudominutum*). The number of assembled genes per transcriptome averaged 48,700, the number of predicted ORFs averaged 41,300, the contig N50 statistic averaged 1,515 bp, mean transcript length per transcriptome averaged 1,078 bp, and annotation success averaged 46.5%. These values are in agreement with previously published Clade B *Symbiodinium* transcriptomes (Bayer et al. 2012; Baumgarten et al. 2013; Shoguchi et al. 2013).

After uploading reference assemblies into IPA software, we identified 19 relevant canonical pathways with significant gene set representation in all species, including fatty acid

**Table 1**Sequencing Statistics for the Ten Strains (A) and Transcriptome Assembly Statistics for the Four Species (B) of Clade B *Symbiodinium*

A								
Species	Strain	Radiation		Ecology		Total Read Count (M)	Remaining After QC (%)	
<i>Symbiodinium minutum</i>	mac703	Pleistocene (B1)		Host-specialized		10.9	89.03	
<i>S. minutum</i>	Mf1.05b	Pleistocene (B1)		Host-specialized		19.3	88.40	
<i>S. minutum</i>	rt002	Pleistocene (B1)		Host-specialized		12.4	88.00	
<i>S. minutum</i>	rt351	Pleistocene (B1)		Host-specialized		8.7	88.68	
<i>Symbiodinium psymophilum</i>	HIAp	Pliocene (B19)		Host-specialized		13.4	88.14	
<i>S. psymophilum</i>	Mf10.14b.02	Pliocene (B19)		Host-specialized		11.1	88.63	
<i>S. psymophilum</i>	PurPflex	Pliocene (B19)		Host-specialized		11.7	88.55	
<i>S. psymophilum</i>	rt141	Pliocene (B19)		Host-specialized		19.5	88.47	
<i>Symbiodinium pseudominutum</i>	rt146	Pleistocene (B1)		Ecologically cryptic		23.7	88.83	
<i>Symbiodinium aenigmaticum</i>	mac04-487	Pliocene (B19)		Ecologically cryptic		11.9	88.10	

B								
Species	Assembly Length (Mbp)	Gene Count	Predicted ORF Count	Genes Annotated (%)	Longest Gene Length (bp)	Mean Gene Length (bp)	N50 (bp)	GC Content (%)
<i>S. minutum</i>	57.2	51,199	42,929	47.3	37,483	1,118	1,579	51.33
<i>S. psymophilum</i>	57.2	50,745	42,740	47.7	31,367	1,128	1,618	51.37
<i>S. pseudominutum</i>	51.3	47,411	40,716	46	31,393	1,081	1,508	51.51
<i>S. aenigmaticum</i>	44.6	45,343	38,923	44.9	24,202	984	1,355	51.39

NOTE.—The *S. minutum* and *S. psymophilum* assemblies in (B) are composited from the reads of all respective strains listed in (A).

beta-oxidation, nitric oxide signaling, oxidative stress response, cell cycle control, RNA processes, and protein ubiquitination (supplementary fig. S1, Supplementary Material online). We compared the ratio of genes observed to total associated genes per pathway among transcriptomes; each pathway was evenly represented in each Clade B *Symbiodinium* species. The four species were also roughly equivalent in terms of their proportions of microsatellite repeat motifs (supplementary fig. S2, Supplementary Material online).


### Between-Species Expression Differences

In order to compare gene expression between *Symbiodinium* species, we identified orthologs via a reciprocal BLAST approach on predicted ORFs. All species shared a total of 19,359 orthologs after filtering out paralogs and low quality matches (average pairwise ortholog count: 27,784; supplementary table S1, Supplementary Material online). We then scaled expression values relative to the lowest expressed gene in the data set (scaled FPKM) to allow for statistical comparison with the R package edgeR. We detected a total of 452 significant DEGs in pairwise species comparisons using expression dispersion estimates derived from all samples across Clade B (fig. 1a). Of these DEGs, 184 could be annotated. The distribution ranged from a low of 45 DEGs between *S. pseudominutum* and *S. psymophilum* to a high of 256 DEGs between *S. minutum* and *S. psymophilum* (fig. 1b). We visualized any annotated gene that was differentially expressed in

at least one species with a heatmap to show relative expression patterns for all samples (fig. 2). A full list of annotations, expression values, and DEG list memberships can be found in supplementary table S3, Supplementary Material online.

We also visualized the ten samples in an MDS plot, which shows spatial clustering based on similarity in gene expression values (fig. 3). When we clustered samples based on the expression of 18,907 genes that were not significantly differentially expressed in species-level contrasts, 8 of the 10 samples spanning three species grouped closely, with only the *S. minutum* strain mac703 and *S. aenigmaticum* strain mac04-487 separating from the others (fig. 3a). When we also included the 452 DEGs, species began to resolve, with one cluster consisting mostly of *S. minutum*, one of mostly *S. psymophilum*, and the single *S. aenigmaticum* remaining unique (fig. 3b). All species segregated well when only the DEGs were considered (fig. 3c).

We subsequently conducted a GO term enrichment analysis on DEGs in order to assess which pathways were differentially represented (supplementary table S2, Supplementary Material online). The *S. psymophilum*–*S. pseudominutum* contrast was enriched for processes including photosynthesis, response to cold, and transmembrane transport. The *S. pseudominutum*–*S. minutum* contrast was enriched for photosynthesis and apoptosis. The *S. aenigmaticum*–*S. pseudominutum* contrast was enriched for photosynthesis and heat acclimation. The *S. psymophilum*–*S. aenigmaticum* contrast was enriched for photosynthesis and mitosis. The *S.*



**FIG. 1.**—(A) Phylogenetic relationships and ecologies for the Clade B *Symbiodinium* species used in this experiment. The maximum parsimony tree was generated based on microsatellite Sym15 flanker region data from Parkinson, Coffroth, et al. (2015) and the methodologies described therein. (B) The numbers of differentially expressed genes (DEGs) between Clade B *Symbiodinium* species. Counts are placed on the lines connecting the two species being contrasted. Line thickness is scaled by the number of DEGs. Also depicted are the numbers ( $n$ ) of cultured strains (clonal cell lines) included per species. Counts below the diagram show the total number of genes differentially expressed in at least one species and the total number of comparable orthologs across all species.

*aenigmaticum*–*S. minutum* contrast was enriched for stress response. The *S. psygmophilum*–*S. minutum* contrast was enriched for photosynthesis, phagocytosis, and cell signaling.

### Within-Species Expression Differences

To understand the extent of gene expression differences among individuals within species, we first quantified expression variability by estimating the BCV and subsequently performed pairwise comparisons of the four distinct strains of *S. minutum* or *S. psygmophilum* using a fixed BCV that was more conservative (lower) than the original BCV estimate (Chen et al. 2014). We identified many pairwise expression differences among strains, ranging from 61 to 404 DEGs for *S. minutum* (fig. 4a) and 82 to 293 DEGs for *S. psygmophilum* (fig. 4b).

To further illustrate differences between pairs of strains for *S. minutum* and *S. psygmophilum*, we calculated an expression–strain distance matrix based on Euclidean distances between all pairwise strain comparisons using scaled FPKM values to assess variance (fig. 4c). Technical variation would be expected to be equally distributed across all samples. The distance matrix revealed a nonrandom distribution of variation in gene expression between pairs of strains for a given species. Also, the distance matrix showed that pairwise distances between strains from different species exceeded any within-species variation, and that both species exhibited distinct variance distributions among member strains.


Finally, we generated heatmaps to show the subset of all annotated genes differentially expressed in at least one individual among the four *S. minutum* strains (fig. 5a) and among the four *S. psygmophilum* strains (fig. 5b). In *S. minutum*,

DEGs between strains were most highly enriched for the processes of malonyl-CoA biosynthesis, protein polymerization, long-chain fatty acid biosynthesis and metabolism, microtubule and nuclear envelope organization, GTP catabolism, and mitosis regulation (supplementary table S2, Supplementary Material online). In *S. psygmophilum*, DEGs between strains were most highly enriched for the processes of DNA replication and biosynthesis, sulfate assimilation and hydrogen sulfide biosynthesis, and microtubule organization (supplementary table S2, Supplementary Material online). A full list of annotations, expression values, and DEG list memberships can be found in supplementary table S3, Supplementary Material online.

### Discussion

Fixed differences in gene expression ultimately influence the phenotypic variation available for selection to act upon. We anticipated that a comparative analysis of *Symbiodinium* spp. transcription would improve our understanding of adaptation and speciation among microeukaryotes. Indeed, we found that despite an overall similarity in gene content and expression among Clade B species with distinct ecologies, all cultures exhibited lineage-specific expression differences diagnostic for each species. Overrepresentation of photosynthesis-related gene expression variation among species likely reflects adaptation to unique light regimes over evolutionary time. Extensive disparity in the expression of fatty acid metabolism genes among strains within species may translate into differences in membrane composition, thermal tolerance, energy reserves, and growth rates. These differences may play a





**Fig. 3.**—Multidimensional scaling plots depicting sample clustering along the primary and secondary leading log<sub>2</sub>-fold change (LFC) axes using the following expression values: (A) Only non-DEGs (18,907), (B) non-DEGs and DEGs together (19,359), and (C) only DEGs (452). No border = host-specialized; border = ecologically cryptic; circles = Pleistocene (B1) radiation, squares = Pliocene (B19) radiation.

crucial role in coral–dinoflagellate symbiosis ecology and evolution. By examining the stable-state transcriptomes of cultures reared independently of their hosts under identical environmental conditions, we infer that these differences stem from genotypic rather than environmental factors. Our efforts reinforce the utility of comparative transcriptomics for studying speciation and functional variation in dinoflagellates and other nonmodel organisms (Chelaifa et al. 2010; Chapman et al. 2013; Gifford et al. 2013).

#### Partitioning the Variation in Gene Expression

When comparing multiple species, expression differences can be attributed to 1) technical variation, 2) within-species variation, and 3) among-species variation, with the proportion of variable genes expected to increase from 1) to 2) to 3) (Whitehead and Crawford 2006). Our results matched this general trend. Technical variation was inferred to be low based on the agreement between our data and transcriptome statistics from other studies that included the same *S. minutum* Mf1.05b strain, the high mapping success achieved between our Mf1.05b reads and the draft genome derived from the same strain (73%), and the nonrandom distribution of expression differences among species (fig. 4c). The percentage of orthologous genes differentially expressed within species (1.54% for *S. minutum* and 1.18% for *S. psygmophilum*; fig. 4) was roughly half that found between species (2.33%; fig. 2b). Overall, DEGs make up a small proportion of the entire transcriptome, as has been found before for *Symbiodinium* and other dinoflagellates (Baumgarten et al. 2013; Barshis et al. 2014; Xiang et al. 2015).

#### Between-Species Variation


By comparing closely related species within a clade, we greatly expanded our comparative power to determine what genetic

changes underlie speciation among *Symbiodinium*. We were able to identify at least four times as many orthologs shared between Clade B species as has been possible using similar methods to compare species across separate clades (Ladner et al. 2012; Barshis et al. 2014; Rosic et al. 2014). Genetic divergence between clades is massive (Rowan and Powers 1992), and thus comparisons among species within clades reveal finer-scale differences likely to be important in physiological and ecological processes. Overall, stable-state gene expression was similar among Clade B *Symbiodinium*. Of the nearly 20,000 orthologs shared by *S. aenigmaticum*, *S. minutum*, *S. pseudominutum*, and *S. psygmophilum*, only 452 (2.3%) were differentially expressed between species. Thus a substantial portion of the transcriptome maintains relatively constant expression levels across members of Clade B. This result mirrors similar studies in other systems such as flowering plants where only a small proportion of interspecific orthologs were differentially expressed (Chapman et al. 2013).

The species comparison with the greatest number of DEGs was *S. minutum* versus *S. psygmophilum* (fig. 1b), which fit expectations for several reasons. First, our replication scheme (four strains per species) may have enhanced our ability to detect fixed differences between these species' transcriptomes (though this is unlikely; see Materials and Methods). Second, both species associate with different hosts and likely diverged in part due to coevolutionary constraints imposed by those hosts, whereas ecologically cryptic species may not have faced the same constraints. Third, they are from phylogenetically divergent lineages. Finally, *S. minutum* is warm water adapted, while *S. psygmophilum* is cold water adapted (Thornhill et al. 2008), likely contributing to expression differences.


Interestingly, the contrasts with the second- and third-most abundant DEG counts both involved *S. aenigmaticum* (fig. 1b), a very distinct species from the Pliocene radiation and

**A** *Symbiodinium minutum*



659 DEGs / 42,929 genes (1.54%)

**B** *Symbiodinium psugmophilum*



506 DEGs / 42,740 genes (1.18%)

**C**

		<i>S. minutum</i>				<i>S. psugmophilum</i>			
		rt351	rt002	mac703	Mf1.05b	rt141	Mf10.14b.02	HIAp	PurPFlex
<i>S. minutum</i>	rt351								
	rt002	3.6							
	mac703	4.5	4.7						
	Mf1.05b	1.7	3.6	5.1					
<i>S. psugmophilum</i>	rt141	10.2	9.3	6.9	10.6				
	Mf10.14b.02	10.1	8.9	6.7	10.3	1.4			
	HIAp	9.7	8.9	5.7	10.2	2.3	2.3		
	PurPFlex	10.2	9.5	6.5	10.6	1.4	2.1	1.5	


**Fig. 4.**—The numbers of DEGs among strains within (A) *Symbiodinium minutum* and (B) *Symbiodinium psugmophilum*. Counts are placed on the lines connecting the two strains being contrasted. Line thickness is scaled by the number of DEGs. Counts below the diagram show the total number of genes differentially expressed in at least one strain and the total number of comparable genes (ORFs) across all strains. (C) Distances between pairs of strains based on Euclidean distance ( $\times 10^3$ ) of expression estimates for all pairwise comparisons of *S. minutum* and *S. psugmophilum* strains. Boxes are shaded such that light coloration reflects relatively small values and dark coloration reflects relatively large values.

one that appears to have undergone rapid evolution (fig. 1a; LaJeunesse 2005; Parkinson, Coffroth, et al. 2015). The three species pair comparisons with the least number of DEGs all involved *S. pseudominutum* (fig. 1b). In fact, this species was roughly equidistant from all other species based on DEG number and MDS position (fig. 3c). Its position might be explained on the one hand by its close evolutionary history with *S. minutum*, and on the other by its cryptic ecology shared with *S. aenigmaticum*. Based on these results, fixed differences in gene expression may not always correspond to phylogenetic similarity.


Multidimensional scaling offered a complementary analysis for visualizing the similarities in expression among all strains

without a priori knowledge of species membership (fig. 3). By restricting the data set to only non-DEGs, almost all replicates from all species (8 of 10) clustered together (fig. 3a), matching the expectation that at stable-state these Clade B *Symbiodinium* generally maintain similar expression profiles. When both non-DEGs and DEGs were included in the analysis, each species was mostly resolved, showing that non-DEGs contributed little to either species-level signal or noise (fig. 3b). As expected, when only the DEGs were considered, all species resolved well (fig. 3c). Note however that the distant positioning of *S. aenigmaticum* in all three MDS plots indicates that a large proportion of expression variation for this species is unique.

**A** *Symbiodinium minutum*



**B** *Symbiodinium psygmpophilum*



**FIG. 5.**—Expression heatmaps of annotated DEGs among individual strains (clonal cell lines) within (A) *Symbiodinium minutum* and (B) *Symbiodinium psygmpophilum*. Colors are scaled to the minimum (purple) and maximum (orange) expression value per gene. Any gene that is differentially expressed in at least one strain is included.

Downloaded from <http://gbe.oxfordjournals.org/> by guest on March 18, 2016

In addition to pairwise comparisons, we also contrasted groups of replicate species by lineage (2 species from the Pleistocene radiation vs. 2 Pliocene radiation species) or by ecology (2 host-specialized species vs. 2 ecologically cryptic species). The Pleistocene—Pliocene contrast was equivalent to the *S. minutum*–*S. psugmophilum* comparison in terms of identity of DEGs, meaning that the species contrast either captured all the differences between major lineages, or that adding just one more strain to each group did not affect expression variation sufficiently to alter our detection of DEGs, even though the strain belonged to a different species. Similarly, the ‘host-specialized’—‘ecologically cryptic’ contrast only recovered four unique genes that had not been identified in any of the species contrasts. The probable identity of only one of these DEGs was determined (a general mRNA splicing factor). These results indicate that differential expression of a particular set of genes does not necessarily explain shared ecological attributes of phylogenetically distinct species.

#### Photosynthesis Gene Expression Differences between Species

Expression differences among closely related species were consistently enriched for photosynthesis genes ([supplementary table S2, Supplementary Material online](#)). Here, overrepresentation of plastid genes cannot be attributed to light intensity differences because all cultures were reared under identical light conditions. In fact, although we might expect these genes to be regulated by light intensity in *Symbiodinium* as they are in other photosynthetic organisms (Escoubas et al. 1995; Pfannschmidt 2003), only minor (or no) changes in photosynthesis-related gene expression have been detected in cultures exposed to varying light levels (McGinley et al. 2013; Xiang et al. 2015). Thus, we conclude that different species evolved unique expression levels among photosynthesis-related genes. These differences may relate to inherent variation in the circadian rhythm among species (Van Dolah et al. 2007; Sorek and Levy 2012) or, more likely, to functional variation in photosynthesis biochemistry. For example, during heat stress, thermally sensitive *Symbiodinium* taxa suffer physiological disruption of PSII photochemistry (Warner et al. 1999; Robison and Warner 2006) and associated downregulation of core photosynthesis genes (McGinley et al. 2012), whereas thermally tolerant species do not. The maintenance of distinct expression patterns at key genes may underlie the capacity for certain *Symbiodinium* species to occupy distinct niches, as has been demonstrated for three diatom species in the genus *Pseudonitzschia* (Di Dato et al. 2015).

#### Evolutionary Significance of Gene Expression Variation

In biogeographic surveys of marine mutualisms, depth and latitude (correlates of light availability) are often primary factors explaining the distribution of *Symbiodinium* diversity

(Rowan and Knowlton 1995; LaJeunesse et al. 2004, 2014; Frade et al. 2008; Finney et al. 2010; Sanders and Palumbi 2011). Thus, light availability represents a main axis of niche differentiation for this group. *Symbiodinium* possess a diverse array of light-harvesting proteins (Boldt et al. 2012), which may be both the cause and consequence of ecological specialization. Many such genes have been transferred to the nuclear genome (Bachvaroff et al. 2004), while others are encoded on plastid minicircles (Zhang et al. 1999; Moore et al. 2003; Barbrook et al. 2014). Minicircles are subject to different transcriptional mechanisms than nuclear encoded genes (Dang and Green 2010), which may also facilitate specialization to different light regimes. Given that a majority of expression variation between divergent species is expected to accumulate neutrally over time (Khaitovich et al. 2005), it is intriguing that expression differences between *Symbiodinium* species are consistently enriched for photosynthesis genes (Baumgarten et al. 2013; Barshis et al. 2014; Rosic et al. 2014; this study). This evidence suggests that species-specific differences in gene expression are functionally important and influenced by natural selection tied to niche diversification.

#### Within-Species Variation

Within each of the two species with four isoclinal cultures, we detected hundreds of DEGs: 659 unique genes among *S. minutum* strains (fig. 4a) and 506 unique genes among *S. psugmophilum* strains (fig. 4b). Interestingly, only four annotated genes differentially expressed among *S. minutum* overlapped with those among *S. psugmophilum*, and enriched categories only overlapped for housekeeping genes which regulate biochemical processes like nucleic acid synthesis and microtubule organization ([supplementary table S2, Supplementary Material online](#)). Thus, transcriptional variation among strains differs from species to species (fig. 5). Furthermore, nonrandom gene expression differences among strains of a given species exist even under identical rearing conditions (fig. 4c), emphasizing that a degree of expression variation among *Symbiodinium* strains is genetically determined and potentially subject to natural selection. Thus, the extent of variation among isoclinal strains may be much greater than previously assumed. Although inter-individual differences are known to play a significant role in symbiosis ecology and evolution in terrestrial systems (Shuster et al. 2006; Whitham et al. 2006; Hughes et al. 2008), such evidence has been lacking for coral–dinoflagellate associations (Parkinson and Baums 2014). Although ~500 of the ~40,000 genes represents a small fraction of the transcriptome, such differences may be important, especially because overall differential expression of genes within a *Symbiodinium* species responding to stress seems low (Barshis et al. 2014; but see Baumgarten et al. 2013).

For example, putative “symbiosis genes” have been identified by comparing symbiotic versus aposymbiotic cnidarian

hosts (Meyer and Weis 2012). The expression levels of similar genes in the symbiont may also play a role in maintaining functional associations. Two such genes varied among *S. minutum* strains: An *ABC transporter* (up to 4.2-fold) and a *glutathione reductase* (up to 9.5-fold). There were also clear differences in the expression of *long chain fatty acid CoA ligase* (up to 12.2-fold), *long chain acyl-CoA synthetase* (up to 8.8-fold), and six *acetyl-CoA carboxylases* (up to 12.5-fold), indicating that certain strains regulate fatty acid metabolism differently. These genes may be related to cell membrane composition, which in turn can affect thermal sensitivity (Tchernov et al. 2004; Diaz-Almeyda et al. 2011). They may also relate to energy storage and nutrient availability, perhaps contributing to different growth rates observed among some of these strains *ex hospite* (Parkinson and Baums 2014). Under environmental change, these functional differences may impact stress tolerance among genotypic host–symbiont combinations in a population (Parkinson and Baums 2014; Parkinson, Banaszak, et al. 2015), partly explaining why some coral colonies of a given species bleach while others do not, even when sharing the same symbiont species (Goulet et al. 2008; Lajeunesse et al. 2010). Similar fine-scale variation has been observed among maize strains with distinct flavonoid content (Casati and Walbot 2003) and among dinoflagellate strains with distinct toxicities (Yang et al. 2010).

## Conclusions

Comparisons among deeply sequenced transcriptomes can reveal the extent and function of molecular variation that is critical to speciation in nonmodel organisms. Such work provides important baselines against which experimentally manipulated samples might be compared and more accurately interpreted. Our data reveal the extent of expression variation that occurs among strains of *Symbiodinium* and emphasizes how natural selection on existing populations may play a critical role in the response of coral–dinoflagellate symbioses to climate change. The genomic resources described here should improve functional investigations into marine symbiosis biology, particularly as model systems continue to be developed (Baumgarten et al. 2015). Future studies should examine the same strains exposed to different stressors (thermal, osmotic, and/or light) in order to characterize the relationship between physiological and gene expression phenotypes. Each strain should also be brought into an experimental host (e.g., the model *Aiptasia* [= *Exaiptasia*]) and observed in symbiosis, which would provide insight into how changes in gene expression work to maintain stable cnidarian–dinoflagellate mutualisms. Our findings underscore that important transcriptional differences exist at different taxonomic ranks among dinoflagellates, from clades to species to strains. Future *Symbiodinium* genomics experiments should be

designed such that clade-level questions incorporate different species to serve as a representative sampling of the clade under study, while species-level questions should incorporate distinct strains to serve as a representative sampling of the species under study. Such designs will improve our understanding of *Symbiodinium* genetic, functional, and phylogenetic diversity.

## Supplementary Material

Supplementary tables S1–S3 and figures S1 and S2 are available at *Genome Biology and Evolution* online (<http://www.gbe.oxfordjournals.org/>).

## Acknowledgments

Mary Alice Coffroth provided cultures, and Maren Ziegler assisted with PRIMER software. This work was supported by the National Science Foundation (DEG-0750756 to J.E.P., OCE-0928764 to T.C.L. and I.B.B.), the Pennsylvania State University, and baseline research funds to CRV by King Abdullah University of Science and Technology (KAUST).

## Literature Cited

- Ahles MD. 1967. Some aspects of the morphology and physiology of *Symbiodinium microadriaticum* [phd dissertation]. [New York]: Fordham University.
- Alexa A, Rahnenfuhrer J. 2010. topGO: enrichment analysis for gene ontology. R package version 2.2.0. <http://bioconductor.uib.no/2.7/bioc/html/topGO.html> Last accessed: January 2014.
- Alexeyenko A, Tamas I, Liu G, Sonnhammer ELL. 2006. Automatic clustering of orthologs and inparalogs shared by multiple proteomes. *Bioinformatics* 22:E9–E15.
- Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search tool. *J Mol Biol*. 215:403–410.
- Bachvaroff TR, Concepcion GT, Rogers CR, Herman EM, Delwiche CF. 2004. Dinoflagellate expressed sequence tag data indicate massive transfer of chloroplast genes to the nuclear genome. *Protist* 155:65–78.
- Barbrook AC, Woolstra CR, Howe CJ. 2014. The chloroplast genome of a *Symbiodinium* sp. clade C3 isolate. *Protist* 165:1–13.
- Barshis DJ, Ladner JT, Oliver J, Palumbi SR. 2014. Lineage specific transcriptional profiles of *Symbiodinium* spp. unaltered by heat stress in a coral host. *Mol Biol Evol*. 31:1343–1352.
- Barshis DJ, et al. 2013. Genomic basis for coral resilience to climate change. *Proc Natl Acad Sci U S A*. 110:1387–1392.
- Baumgarten S, et al. 2013. Integrating microRNA and mRNA expression profiling in *Symbiodinium microadriaticum*, a dinoflagellate symbiont of reef-building corals. *BMC Genomics* 14:704
- Baumgarten S, et al. 2015. The genome of *Aiptasia*, a sea anemone model for coral symbiosis. *Proc Natl Acad Sci U S A*. 112:11893–11898.
- Bayer T, et al. 2012. *Symbiodinium* transcriptomes: genome insights into the dinoflagellate symbionts of reef-building corals. *PLoS One* 7:e5269
- Boldt L, Yellowlees D, Leggat W. 2012. Hyperdiversity of genes encoding integral light-harvesting proteins in the dinoflagellate *Symbiodinium* sp. *PLoS One* 7:e47456
- Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114–2120.

- Casati P, Walbot V. 2003. Gene expression profiling in response to ultraviolet radiation in maize genotypes with varying flavonoid content. *Plant Physiol.* 132:1739–1754.
- Chapman MA, Hiscock SJ, Filatov DA. 2013. Genomic divergence during speciation driven by adaptation to altitude. *Mol Biol Evol.* 30:2553–2567.
- Chelaifa H, Mahe F, Ainouche M. 2010. Transcriptome divergence between the hexaploid salt-marsh sister species *Spartina maritima* and *Spartina alterniflora* (Poaceae). *Mol Ecol.* 19:2050–2063.
- Chen Y, Lun AT, Smyth GK. 2014. Differential expression analysis of complex RNA-seq experiments using edgeR. In: Nettleton D, Datta S, editors. *Statistical analysis of next generation sequencing data*. Springer International Publishing: Switzerland. p. 51–74.
- Clarke K, Gorley R. 2006. *PRIMER v6: user manual/tutorial*. Plymouth (UK): PRIMER-E. 192pp.
- Dang YK, Green BR. 2010. Long transcripts from dinoflagellate chloroplast minicircles suggest “rolling circle” transcription. *J Biol Chem.* 285:5196–5203.
- Diaz-Almeyda E, Thome PE, El Hafidi M, Iglesias-Prieto R. 2011. Differential stability of photosynthetic membranes and fatty acid composition at elevated temperature in *Symbiodinium*. *Coral Reefs* 30:217–225.
- Di Dato V, et al. 2015. Transcriptome sequencing of three *Pseudo-nitzschia* species reveals comparable gene sets and the presence of *Nitric Oxide Synthase* genes in diatoms. *Sci Rep.* 5:12329
- Dimmer EC, et al. 2012. The UniProt-GO Annotation database in 2011. *Nucleic Acids Res.* 40:D565–D570.
- Erdner DL, Anderson DM. 2006. Global transcriptional profiling of the toxic dinoflagellate *Alexandrium fundyense* using massively parallel signature sequencing. *BMC Genomics* 7:88
- Escoubas JM, Lomas M, Laroche J, Falkowski PG. 1995. Light-intensity regulation of *cab* gene: transcription is signaled by the redox state of the plastoquinone pool. *Proc Natl Acad Sci U S A.* 92:10237–10241.
- Finn RD, et al. 2014. Pfam: the protein families database. *Nucleic Acids Res.* 42:D222–D230.
- Finney JC, et al. 2010. The relative significance of host-habitat, depth, and geography on the ecology, endemism, and speciation of coral endosymbionts in the genus *Symbiodinium*. *Microb Ecol.* 60:250–263.
- Fitt WK, Brown BE, Warner ME, Dunne RP. 2001. Coral bleaching: interpretation of thermal tolerance limits and thermal thresholds in tropical corals. *Coral Reefs* 20:51–65.
- Frade PR, De Jongh F, Vermeulen F, Van Bleijswijk J, Bak RPM. 2008. Variation in symbiont distribution between closely related coral species over large depth ranges. *Mol Ecol.* 17:691–703.
- Galardini M, et al. 2011. Exploring the symbiotic pangenome of the nitrogen-fixing bacterium *Sinorhizobium meliloti*. *BMC Genomics* 12:235
- Gifford SM, Sharma S, Booth M, Moran MA. 2013. Expression patterns reveal niche diversification in a marine microbial assemblage. *ISME J.* 7:281–298.
- Gnerre S, et al. 2011. High-quality draft assemblies of mammalian genomes from massively parallel sequence data. *Proc Natl Acad Sci U S A.* 108:1513–1518.
- Gould J. 2015. Gene-E. Available from: <http://www.broadinstitute.org/cancer/software/GENE-E/index.html>. Last accessed: June 2015.
- Goulet TL, LaJeunesse TC, Fabricius KE. 2008. Symbiont specificity and bleaching susceptibility among soft corals in the 1998 Great Barrier Reef mass coral bleaching event. *Mar Biol.* 154:795–804.
- Grabherr MG, et al. 2011. Full-length transcriptome assembly from RNA-seq data without a reference genome. *Nat Biotechnol.* 29:644–652.
- Haas BJ, et al. 2013. *De novo* transcript sequence reconstruction from RNA-seq using the Trinity platform for reference generation and analysis. *Nat Protoc.* 8:1494–1512.
- Hoegh-Guldberg O. 1999. Climate change, coral bleaching and the future of the world's coral reefs. *Mar Freshwater Res.* 50:839–866.
- Hughes AR, Inouye BD, Johnson MTJ, Underwood N, Vellend M. 2008. Ecological consequences of genetic diversity. *Ecol Lett.* 11:609–623.
- Jeong HJ, et al. 2014. Genetics and morphology characterize the dinoflagellate *Symbiodinium voratum*, n. sp., (Dinophyceae) as the sole representative of *Symbiodinium* clade E. *J Eukaryot Microbiol.* 61:75–94.
- Khaitovich P, Paabo S, Weiss G. 2005. Toward a neutral evolutionary model of gene expression. *Genetics* 170:929–939.
- Kofler R, Schlotterer C, Lelley T. 2007. SciRoKo: a new tool for whole genome microsatellite search and investigation. *Bioinformatics* 23:1683–1685.
- Ladner JT, Barshis DJ, Palumbi SR. 2012. Protein evolution in two co-occurring types of *Symbiodinium*: an exploration into the genetic basis of thermal tolerance in *Symbiodinium* clade D. *BMC Evol Biol.* 12:217
- LaJeunesse TC. 2005. “Species” radiations of symbiotic dinoflagellates in the Atlantic and Indo-Pacific since the Miocene-Pliocene transition. *Mol Biol Evol.* 22:570–581.
- LaJeunesse TC, Lee SY, Gil-Agudelo DL, Knowlton N, Jeong HJ. 2015. *Symbiodinium necroappetens* sp. nov. (Dinophyceae): an opportunistic ‘zooxanthella’ found in bleached and diseased tissues of Caribbean reef corals. *Eur J Phycol.* 50:223–238.
- LaJeunesse TC, Parkinson JE, Reimer JD. 2012. A genetics-based description of *Symbiodinium minutum* sp. nov. and *S. psygmophilum* sp. nov. (Dinophyceae), two dinoflagellates symbiotic with cnidaria. *J Phycol.* 48:1380–1391.
- LaJeunesse TC, Thornhill DJ. 2011. Improved resolution of reef-coral endosymbiont (*Symbiodinium*) species diversity, ecology, and evolution through *psbA* non-coding region genotyping. *PLoS One* 6:e29013.
- LaJeunesse TC, et al. 2004. Closely related *Symbiodinium* spp. differ in relative dominance in coral reef host communities across environmental, latitudinal and biogeographic gradients. *Mar Ecol Prog Ser.* 284:147–161.
- LaJeunesse TC, et al. 2010. Host-symbiont recombination versus natural selection in the response of coral-dinoflagellate symbioses to environmental disturbance. *Proc Biol Sci.* 277:2925–2934.
- LaJeunesse TC, et al. 2014. Ecologically differentiated stress tolerant endosymbionts in the dinoflagellate genus *Symbiodinium* Clade D are different species. *Phycologia* 53:305–319.
- Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. *Nat Methods.* 9:357–359.
- Leggat W, Hoegh-Guldberg O, Dove S, Yellowlees D. 2007. Analysis of an EST library from the dinoflagellate (*Symbiodinium* sp.) symbiont of reef-building corals. *J Phycol.* 43:1010–1021.
- Leggat W, Seneca F, et al. 2011. Differential responses of the coral host and their algal symbiont to thermal stress. *PLoS One* 6:e26687
- Leggat W, Yellowlees D, Medina M. 2011. Recent progress in *Symbiodinium* transcriptomics. *J Exp Mar Biol Ecol.* 408:120–125.
- Lehnert EM, Burriesci MS, Pringle JR. 2012. Developing the anemone *Aiptasia* as a tractable model for cnidarian-dinoflagellate symbiosis: the transcriptome of aposymbiotic *A. pallida*. *BMC Genomics* 13:271.
- Lehnert EM, et al. 2014. Extensive differences in gene expression between symbiotic and aposymbiotic cnidarians. *G3 (Bethesda)* 4:277–295.
- Lin SJ. 2011. Genomic understanding of dinoflagellates. *Res Microbiol.* 162:551–569.
- Lin SJ, et al. 2015. The *Symbiodinium kawagutii* genome illuminates dinoflagellate gene expression and coral symbiosis. *Science* 350:691–694.
- McCarthy DJ, Chen Y, Smyth GK. 2012. Differential expression analysis of multifactor RNA-Seq experiments with respect to biological variation. *Nucleic Acids Res.* 40:4288–4297.
- McGinley MP, Suggett DJ, Warner ME. 2013. Transcript patterns of chloroplast-encoded genes in cultured *Symbiodinium* spp. (Dinophyceae): testing the influence of a light shift and diel periodicity. *J Phycol.* 49:709–718.
- McGinley MP, et al. 2012. Transcriptional response of two core photosystem genes in *Symbiodinium* spp. exposed to thermal stress. *PLoS One* 7:e50439

- McLaughlin JJA, Zahl PA. 1959. Axenic zooxanthellae from various invertebrate hosts. *Ann N Y Acad Sci.* 77:55–72.
- Meyer E, Weis VM. 2012. Study of cnidarian-algal symbiosis in the “omics” age. *Biol Bull.* 223:44–65.
- Moberg F, Folke C. 1999. Ecological goods and services of coral reef ecosystems. *Ecol Econ.* 29:215–233.
- Moore RB, Ferguson KM, Loh WKW, Hoegh-Guldberg C, Carter DA. 2003. Highly organized structure in the non-coding region of the *psbA* minicircle from clade C *Symbiodinium*. *Int J Syst Evol Microbiol.* 53:1725–1734.
- Morse D, Milos PM, Roux E, Hastings JW. 1989. Circadian regulation of bioluminescence in *Gonyaulax* involves translational control. *Proc Natl Acad Sci U S A.* 86:172–176.
- Moustafa A, et al. 2010. Transcriptome profiling of a toxic dinoflagellate reveals a gene-rich protist and a potential impact on gene expression due to bacterial presence. *PLoS One* 5:e9688
- Muscatine L. 1990. The role of symbiotic algae in carbon and energy flux in reef corals. In: Dubinsky, editor. *Ecosystems of the world 25: Coral reefs*. New York: Elsevier. p. 75–87.
- Muscatine L, Porter JW. 1977. Reef corals—mutualistic symbioses adapted to nutrient-poor environments. *Bioscience* 27:454–460.
- Österman J, Mousavi SA, Koskinen P, Paulin L, Lindström K. 2015. Genomic features separating ten strains of *Neorhizobium galegae* with different symbiotic phenotypes. *BMC Genomics* 16:348
- Parkinson JE, Banaszak AT, Altman NS, LaJeunesse TC, Baums IB. 2015. Intraspecific diversity among partners drives functional variation in coral symbioses. *Sci Rep.* 5:15667
- Parkinson JE, Baums IB. 2014. The extended phenotypes of marine symbioses: ecological and evolutionary consequences of intraspecific genetic diversity in coral-algal associations. *Front Microbiol.* 5:445
- Parkinson JE, Coffroth MA, LaJeunesse TC. 2015. New species of Clade B *Symbiodinium* (Dinophyceae) from the greater Caribbean belong to different functional guilds: *S. aenigmaticum* sp. nov., *S. antillogorgium* sp. nov., *S. endomadracis* sp. nov., and *S. pseudominutum* sp. nov. *J Phycol.* 51:850–858.
- Pettay DT, LaJeunesse TC. 2007. Microsatellites from clade B *Symbiodinium* spp. specialized for Caribbean corals in the genus *Madracis*. *Mol Ecol Notes.* 7:1271–1274.
- Pfannschmidt T. 2003. Chloroplast redox signals: how photosynthesis controls its own genes. *Trends Plant Sci.* 8:33–41.
- Provasoli L. 1968. Media and prospects for the cultivation of marine algae. In: Watanabe H, Hattori A, editors. *Culture and Collection of Algae. Proceedings of the US-Japan Conference*. Japanese Society for Plant Physiology; Hakone, Japan. p. 63–75.
- Pruitt KD, et al. 2014. RefSeq: an update on mammalian reference sequences. *Nucleic Acids Res.* 42:D756–D763.
- Ranz JM, Machado CA. 2006. Uncovering evolutionary patterns of gene expression using microarrays. *Trends Ecol Evol.* 21:29–37.
- Remm M, Storm CEV, Sonnhammer ELL. 2001. Automatic clustering of orthologs and in-paralogs from pairwise species comparisons. *J Mol Biol.* 314:1041–1052.
- Ribeiro FJ, et al. 2012. Finished bacterial genomes from shotgun sequence data. *Genome Res.* 22:2270–2277.
- Roberts A, Pachter L. 2013. Streaming fragment assignment for real-time analysis of sequencing experiments. *Nat Methods.* 10:71–73.
- Robinson MD, McCarthy DJ, Smyth GK. 2010. edgeR: a bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics* 26:139–140.
- Robison JD, Warner ME. 2006. Differential impacts of photoacclimation and thermal stress on the photobiology of four different phylotypes of *Symbiodinium* (Pyrrophyta). *J Phycol.* 42:568–579.
- Rosic N, et al. 2014. Unfolding the secrets of coral-algal symbiosis. *ISME J.* 9:844–856.
- Rowan R, Knowlton N. 1995. Intraspecific diversity and ecological zonation in coral algal symbiosis. *Proc Natl Acad Sci U S A.* 92:2850–2853.
- Rowan R, Powers DA. 1992. Ribosomal RNA sequences and the diversity of symbiotic dinoflagellates (Zooxanthellae). *Proc Natl Acad Sci U S A.* 89:3639–3643.
- Sanders JG, Palumbi SR. 2011. Populations of *Symbiodinium muscatinei* show strong biogeographic structuring in the intertidal anemone *Anthopleura elegantissima*. *Biol Bull.* 220:199–208.
- Santos SR, Taylor DJ, Coffroth MA. 2001. Genetic comparisons of freshly isolated versus cultured symbiotic dinoflagellates: implications for extrapolating to the intact symbiosis. *J Phycol.* 37:900–912.
- Schoenberg DA, Trench RK. 1980. Genetic variation in *Symbiodinium* (= *Gymnodinium*) *microadriaticum* Freudenthal, and specificity in its symbiosis with marine invertebrates. 1. Isoenzyme and soluble-protein patterns of axenic cultures of *Symbiodinium microadriaticum*. *Proc Biol Sci.* 207:405–427.
- Shoguchi E, et al. 2013. Draft assembly of the *Symbiodinium minutum* nuclear genome reveals dinoflagellate gene structure. *Curr Biol.* 23:1399–1408.
- Shuster SM, Lonsdorf EV, Wimp GM, Bailey JK, Whitham TG. 2006. Community heritability measures the evolutionary consequences of indirect genetic effects on community structure. *Evolution* 60:991–1003.
- Sorek M, Levy O. 2012. Influence of the quantity and quality of light on photosynthetic periodicity in coral endosymbiotic algae. *PLoS One* 7:e43264
- Sunagawa S, et al. 2009. Generation and analysis of transcriptomic resources for a model system on the rise: the sea anemone *Aiptasia pallida* and its dinoflagellate endosymbiont. *BMC Genomics* 10:258
- Tchernov D, et al. 2004. Membrane lipids of symbiotic algae are diagnostic of sensitivity to thermal bleaching in corals. *Proc Natl Acad Sci U S A.* 101:13531–13535.
- Thornhill DJ, Kemp DW, Bruns BU, Fitt WK, Schmidt GW. 2008. Correspondence between cold tolerance and temperate biogeography in a Western Atlantic *Symbiodinium* (Dinophyta) lineage. *J Phycol.* 44:1126–1135.
- UniProt Consortium 2013. Update on activities at the Universal Protein Resource (UniProt) in 2013. *Nucleic Acids Res.* 41:D43–D47.
- Van Dolah FM, et al. 2007. Microarray analysis of diurnal- and circadian-regulated genes in the Florida red-tide dinoflagellate *Karenia brevis* (Dinophyceae). *J Phycol.* 43:741–752.
- Voolstra C, Tautz D, Farbrother P, Eichinger L, Harr B. 2007. Contrasting evolution of expression differences in the testis between species and subspecies of the house mouse. *Genome Res.* 17:42–49.
- Voolstra CR, et al. 2009. The host transcriptome remains unaltered during the establishment of coral-algal symbioses. *Mol Ecol.* 18:1823–1833.
- Warner ME, Fitt WK, Schmidt GW. 1999. Damage to photosystem II in symbiotic dinoflagellates: a determinant of coral bleaching. *Proc Natl Acad Sci U S A.* 96:8007–8012.
- Weis VM, Davy SK, Hoegh-Guldberg O, Rodriguez-Lanetty M, Pringle JR. 2008. Cell biology in model systems as the key to understanding corals. *Trends Ecol Evol.* 23:369–376.
- Whitehead A, Crawford DL. 2006. Variation within and among species in gene expression: raw material for evolution. *Mol Ecol.* 15:1197–1211.
- Whitham TG, et al. 2006. A framework for community and ecosystem genetics: from genes to ecosystems. *Nat Rev Genet.* 7:510–523.
- Wittkopp PJ, Haerum BK, Clark AG. 2008. Regulatory changes underlying expression differences within and between *Drosophila* species. *Nat Genet.* 40:346–350.

- Wolf JBW, Lindell J, Backstrom N. 2010. Speciation genetics: current status and evolving approaches. *Philos Trans R Soc B Biol Sci.* 365:1717–1733.
- Xiang T, Nelson W, Rodriguez J, Tolleter D, Grossman AR. 2015. *Symbiodinium* transcriptome and global responses of cells to immediate changes in light intensity when grown under autotrophic or mixotrophic conditions. *Plant J.* 82:67–80.
- Xiang TT, Hambleton EA, DeNofrio JC, Pringle JR, Grossman AR. 2013. Isolation of clonal axenic strains of the symbiotic dinoflagellate *Symbiodinium* and their growth and host specificity. *J Phycol.* 49:447–458.
- Yang I, et al. 2010. Comparative gene expression in toxic versus non-toxic strains of the marine dinoflagellate *Alexandrium minutum*. *BMC Genomics* 11:248
- Yellowlees D, Rees TAV, Leggat W. 2008. Metabolic interactions between algal symbionts and invertebrate hosts. *Plant Cell Environ.* 31:679–694.
- Zhang Z, Green B, Cavalier-Smith T. 1999. Single gene circles in dinoflagellate chloroplast genomes. *Nature* 400:155–159.

Associate editor: Eric Bapteste